

Pacific Islands - Online Climate Outlook Forum (OCOF) No. 119

Country Name: Samoa

TABLE 1: Monthly Rainfall

Station (include data period)	July 2017						
	May 2017 Total	June 2017 Total	Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking
Afiamalu	1007.9	354.2	120.5	145.5	257.7	207.6	18/66
Nafanua	951.5	206.9	142.8	84.6	168.6	117.7	28/46
Apia	882.2	162.4	104.1	71.6	129.5	93.3	75/128
Faleolo	420.4	145.0	26.7	53.2	103.4	80.9	8/57

TABLE 2: Three-monthly Rainfall May to July 2017

[Please note that the data used in this verification should be sourced from table 3 of OCOF #115]

Station	Three-month Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking	Forecast probs.* (include LEPS)	Verification* (Consistent, Near-consistent Inconsistent?)
Afiamalu	1482.6	625.1	832.5	732.4	61/63	29/36/35 (1.0)	Near consistent
Nafanua	1301.2	437.9	594.1	523.6	45/45	32/34/34 (-2.6)	Near-Consistent
Apia	1148.7	333.0	481.2	410.2	128/128	33/34/33 (-2.3)	Near consistent
Faleolo	592.1	273.1	374.3	311.3	55/55	32/31/37 (0.9)	Consistent

Period: *below normal/normal/above normal

Predictors and Period used for May to July 2017 Outlooks (refer to OCOF #115):
Nino 3.4 indices from February to March 2017.

* Forecast is consistent when observed and predicted (tercile with the highest probability) categories coincide (are in the same tercile).

Forecast is near-consistent when observed and predicted (tercile with the highest probability) differ by only one category (i.e. terciles 1 and 2 or terciles 2 and 3).

Forecast is inconsistent when observed and predicted (tercile with the highest probability) differ by two categories (i.e. terciles 1 and 3).

**TABLE 3: Seasonal Climate Outlooks using SCOPIC for
September to November 2017**

Predictors and Period used: Nino 3.4 values of June to July 2017

Station	Below Median (prob)	Median Rainfall (mm)	Above Median (prob)		LEPS	Hit-rate
Afiamalu	56	909.9	44		3.5%	60.7%
Nafanua	54	675.1	46		2.1%	56.8%
Apia	57	593.4	43		5.9 %	61.2%
Faleolo	51	468.3	49		-1.1%	54.7%

Station	Below Normal (prob)	33%ile rainfall (mm)	Normal (prob)	66%ile rainfall (mm)	Above Normal (prob)	LEPS	Hit-rate
Afiamalu	38	777.8	34	1094.1	28	3.5%	42.9%
Nafanua	38	598.6	32	791.7	30	3.0%	38.6%
Apia	38	493.0	33	682.1	29	1.8%	22.4%
Faleolo	36	401.8	33	518.2	31	0.1%	28.3%

**TABLE 4: Seasonal Climate Outlooks using POAMA2 for
September to November 2017**

Station	Lower Tercile (prob)	33%ile rainfall (mm)	Middle Tercile (prob)	66%ile rainfall (mm)	Upper Tercile (prob)		
Apia	15	497	6	679	79		
Faleolo	15	300	6	486	79		

Summary Statements

Rainfall for July 2017:

'Below normal' rainfall was recorded at Afiamalu and Faleolo station whereas Apia and Nafanua registered 'normal'.

Accumulated rainfall for May to July 2017, including outlook verification:

'Above normal' rainfall for all stations. New records for Apia, Afiamalu and Faleolo as they recorded highest ranking accumulated rainfall in this period.

Apia, Nafanua and Afiamalu were 'near consistent' with the outlook while Faleolo recorded 'consistent'.

Outlooks for August to October 2017:

1. SCOPIC:

The outlook offers little guidance for Afiamalu, Apia and Faleolo as the chances of '**below normal**, '**normal**' and '**above normal**' are similar.

'**Below normal**' rainfall is anticipated for Nafanua with '**normal**' the next most likely outcome.

The model confidence is low

2. POAMA:

'**Above normal**' rainfall is the favoured outcome for Apia and Faleolo stations.

NB: The X LEPS % score has been categorised as follows:

Very Low: $X < 0.0$

Low: $0 \leq X < 5$

Moderate $5 \leq X < 10$

Good: $10 \leq X < 15$

High: $15 \leq X < 25$

Very High: $25 \leq X < 35$

Exceptional: $X \geq 35$