Pacific Islands - Online Climate Outlook Forum (OCOF) No. 67

Country Name: TONGA

TABLE 1: Monthly Rainfall

Station (include data period)			March 2013						
	January 2013 Total	February 2013 Total	Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking		
Northern Division									
Niuafoóu	362.6	314.0	450.4	213.0	348.0	268.4	34/43		
Niuatoputapu	573.0	210.2	116.8	187.0	322.7	241.5	8/67		
Central Division									
Vavaú	316.4	201.6	311.3	227.3	366.7	301.0	35/67		
Haápai	282.6	271.4	335.4	179.0	313.0	233.5	51/67		
Southern Division									
Nukuálofa	301.3	287.8	531.2	168.3	252.6	209.0	69/69		
Fuaámotu	335.2	550.5	445.0	145.3	213.3	173.0	34/34		

TABLE 2: Three-monthly Rainfall

January 2013 to March 2013

[Please note that the data used in this verification should be sourced from table 3 of OCOF #62]

Station	Three-month Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking	Forecast probs.* (include LEPS)	Verification* (Consistent, Near-consistent Inconsistent?		
Northern Division									
Niuafoóu	1127.0	753.0	955.2	893.5	30/43	21,24,55 (9.1)	Consistent		
Niuatoputapu	900.0	650.0	939.7	810.0	41/67	44 ,36,20 (12.9)	Near consistent		
Central Division									
Vavaú	829.3	687.3	946.7	858.5	31/67	40 ,29,31 (9.9)	Near Consistent		
Haápai	889.4	538.4	764.0	651.0	56/67	27, 37 ,36 (26.2)	Near Consistent		
Southern Division									
Nukuálofa	1120.3	485.3	807.3	633.0	67/69	27 ,58 ,15 (15.9)	Near Consistent		
Fuaámotu	1330.7	455.3	770.3	584.0	34/34	24, 51 ,25 (26.2)	Near Consistent		

Period:*below normal/normal/above normal

<u>Predictors and Period used for January 2013 to March 2013 Outlooks (refer to OCOF #62):</u> **Preditors – SSTa's I@9 Period; Sept – November with 1 month lead.**

^{*}Forecast is <u>consistent</u> when observed and predicted (tercile with the highest probability) categories coincide (are in the same tercile).

Forecast is <u>near-consistent</u> when observed and predicted (tercile with the highest probability) differ by only one category (i.e. terciles 1 and 2 or terciles 2 and 3).

Forecast is <u>inconsistent</u> when observed and predicted (tercile with the highest probability) differ by two categories (i.e. terciles 1 and 3).

TABLE 3: Seasonal Climate Outlooks using SCOPIC for May to July 2013

Predictors and Period used: SSTa's 1@9 – January 2013 to March 2013

Station	Below Normal (prob)	33%ile rainfall (mm)	Normal (prob)	66%ile rainfall (mm)	Above Normal (prob)	LEPS	Hit-rate
Northern Division Niuafoóu Niuatoputapu Central Division Vavaú Haápai	32 28 31 32	373.7 321.3 300.0 208.3	32 41 32 33	476.9 405.0 402.0 307.7	36 31 37 35	-2.6 2.3 11.4 5.3	23.7 39.3 42.2 33.3
Southern Division Nukuálofa Fuaámotu	22 25	237.3 273.6	39 37	343.0 441.9	39 38	15.6 10.9	32.8 34.4

TABLE 4: Seasonal Climate Outlooks using POAMA2 for May to July 2013

Station	Lower Tercile (prob)	33%ile rainfall (mm)	Middle Tercile (prob)	66%ile rainfall (mm)	Upper Tercile (prob)	
Nukuálofa						 - - -

Summary Statements

Rainfall for March 2013:

Northern Division: Above normal for Niuafo'ou. Below normal for Niuatoputapu.

Central Division: Normal to above normal.

Southern Division: Above Normal.

Accumulated rainfall for December 2012–February 2013, including outlook verification:

Northern Division: Normal to Above Normal. Forecast was Near Consistent to Consistent.

Central Division: Normal to Above Normal. Forecast was Near Consistent

Southern Division: Above Normal. Forecast was Near Consistent.

Outlooks for April-June 2013:

1. SCOPIC:

Northern Division: Normal to above normal.

Central Division: Normal to above normal.

Southern Division: Above Normal.

2. POAMA: Outlook

NB: The X LEPS % score has been categorised as follows:

 $\mbox{Very Low: } X < 0.0 \qquad \qquad \mbox{Low: } 0 \le X < 5 \qquad \qquad \mbox{Moderate } 5 \le X < 10 \qquad \qquad \mbox{Good: } 10 \le X < 15 \qquad \mbox{High: } 15 \le X < 25$

Very High: $25 \le X < 35$ Exceptional: $X \ge 35$