

Pacific Islands - Online Climate Outlook Forum (OCOF) No. 85

Country Name: TONGA

TABLE 1: Monthly Rainfall

Station (include data period)	September 2014						
	July 2014 Total	August 2014 Total	September 2014 Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking
Northern Division							
Niuafoóu	11.7	37.1	102.4	82.3	155.7	131.0	19/44
Niuaotoputapu	125.2	148	16.0	53.0	131.4	100.5	5/68
Central Division							
Vavaú	54.9	42.4	33.9	81.0	167.0	109.0	6/68
Haápai	47.2	34.6	1.1	51.0	144.0	91.0	1/68
Southern Division							
Nukuálofa	157.7	15.3	32.9	75.7	144.7	103.0	6/70
Fuaámotu	121.3	13.9	19.0	52.0	153.0	105.0	1/35

TABLE 2: Three-monthly Rainfall

July to September 2014

[Please note that the data used in this verification should be sourced from table 3 of OCOF #81]

Predictors and Period used: NINO3.4 SST – Mar to May 2014

Station	Three-month Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking	Forecast probs.* (include LEPS)	Verification* (Consistent, Near-consistent, Inconsistent?)
Northern Division							
Niuafoóu	151.2	269.0	422.6	338.5	3/44	29, 32, 39 (-4.7)	In consistent
Niuaotoputapu	289.2	206.0	365.0	262.5	35/68	Missing data in Niuaotoputapu was due on the month of May	
Central Division							
Vavaú	131.2	254.0	399.0	340	4/68	24, 35, 41 (-1.1)	Near consistent
Haápai	82.9	237.5	365.0	294.6	1/68	34, 22, 44 (-0.6)	Near consistent
Southern Division							
Nukuálofa	205.9	276.3	383.3	323.7	16/70	38, 18, 44 (-2.7)	Near consistent
Fuaámotu	154.2	298.0	423.0	374.0	3/35	41, 18, 41 (-5.3)	In consistent

Period: *below normal/normal/above normal

* Forecast is consistent when observed and predicted (tercile with the highest probability) categories coincide (are in the same tercile).

Forecast is near-consistent when observed and predicted (tercile with the highest probability) differ by only one category (i.e. terciles 1 and 2 or terciles 2 and 3).

Forecast is inconsistent when observed and predicted (tercile with the highest probability) differ by two categories (i.e. terciles 1 and 3).

TABLE 3: Seasonal Climate Outlooks using SCOPIC for November 2014 to January 2015

Predictors and Period used: NINO 3.4 SST Anomalies– July to September 2014

Station	Below Median (prob)	Median Rainfall (mm)	Above Median (prob)		LEPS	Hit-rate
Northern Division						
Niuafo'ou	52	825.6	48		1.6	58.3
Niuaatoputapu	55	755	45		14	71.2
Central Division						
Vava'u	58	655	42		27.2	76.2
Ha'apai	57	459	43		22.1	73.4
Southern Division						
Nuku'alofoa	57	439.5	43		20.8	76.6
Fua'amotu	51	467.5	49		13.3	70.6

Station	Below Normal (prob)	33%ile rainfall (mm)	Normal (prob)	66%ile rainfall (mm)	Above Normal (prob)	LEPS	Hit-rate
Northern Division							
Niuafo'ou	34	710	33	1005.9	33	-0.4	47.2
Niuaatoputapu	37	622	37	867	26	14.4	47.5
Central Division							
Vava'u	37	492	39	818	24	21.3	50.8
Ha'apai	36	324	38	571	26	23.8	59.4
Southern Division							
Nuku'alofoa	35	342	40	568	25	28.6	54.7
Fua'amotu	33	392	38	616	29	14.2	50

TABLE 4: Seasonal Climate Outlooks using POAMA2 for November 2014 to January 2015

Station	Lower Tercile (prob)	33%ile rainfall (mm)	Middle Tercile (prob)	66%ile rainfall (mm)	Upper Tercile (prob)		
Nuku'alofoa	24.	229	55	380	21		

Summary Statements

Rainfall for September 2014:

Northern Division: Niuafu'ou - Below normal

Niutoputapu - Normal

Central Division: Below normal

Southern Division: Below normal

Accumulated rainfall for July-September 2014, including outlook verification:

Northern Division: Niuafu'ou - Below normal, outlook was inconsistent

Niutoputapu - Normal

Central Division: Below normal, outlook was near consistent

Southern Division: Nuku'alofa - Below normal, outlook was near consistent.

Fua'amotu - Below normal, outlook was inconsistent

Outlooks for November 2014 - January 2015:

1. SCOPIC:

The SCOPIC seasonal rainfall outlook for November 2014 to January 2015 shows:

- At Niuafu'ou in the Northern Division the chances of above-normal, normal and below-normal rainfall are similar. At Niutoputapu there is an equal likelihood of below normal and normal rainfall.
- In the Central Division there is an equal likelihood of below normal and normal rainfall.
- At Nuku'alofa in the Southern Division, the most likely outcome is normal rainfall, with below normal the next most likely. At Fua'amotu the chances of above-normal, normal and below-normal rainfall are similar.

2. POAMA:

- POAMA seasonal Outlook for Nuku'alofa for November to January 2015 shows the most likely outcome is Normal

NB: The X LEPS % score has been categorised as follows:

Very Low: $X < 0.0$

Low: $0 \leq X < 5$

Moderate $5 \leq X < 10$

Good: $10 \leq X < 15$

High: $15 \leq X < 25$

Very High: $25 \leq X < 35$

Exceptional: $X \geq 35$