

Pacific Islands - Online Climate Outlook Forum (OCOF) No. 91

Country Name: Samoa

TABLE 1: Monthly Rainfall

Station (include data period)	March 2015						
	January 2015 Total	February 2015 Total	Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking
Afiamalu	626.4	534.7	552.6	392.9	593	501.5	39/63
Nafanua	437.4	382.6	320.0	312.3	410.2	358.4	16/43
Apia	370.2	348.6	314.4	273.3	377	309.2	65/126
Faleolo	188.3	238.1	256.0	180.5	243.7	213.6	38/54

TABLE 2: Three-monthly Rainfall January 2015 to March 2015

[Please note that the data used in this verification should be sourced from table 3 of OCOF #87]

Station	Three-month Total	33%tile Rainfall (mm)	67%tile Rainfall (mm)	Median Rainfall (mm)	Ranking	Forecast probs.* (include LEPS)	Verification* (Consistent, Near-consistent Inconsistent?)
Afiamalu	1713.7	1562.7	1981.8	1775.1	28/61	38/38/24 (4.5)	Near Consistent
Nafanua	1140.0	1097.2	1609.2	1303.7	17/41	45/34/22 (16.9)	Near consistent
Apia	1033.2	917.0	1171.4	1056.8	45/126	44/31/25 (13.3)	Near consistent
Faleolo	682.4	758.3	897.3	856.3	18/54	38/26/36 (-3.3)	Consistent

Period: *below normal/normal/above normal

Predictors and Period used for January to March 2015 Outlooks (refer to OCOF #87):
Nino 3.4 value from September to November 2014.

* Forecast is consistent when observed and predicted (tercile with the highest probability) categories coincide (are in the same tercile).

Forecast is near-consistent when observed and predicted (tercile with the highest probability) differ by only one category (i.e. terciles 1 and 2 or terciles 2 and 3).

Forecast is inconsistent when observed and predicted (tercile with the highest probability) differ by two categories (i.e. terciles 1 and 3).

**TABLE 3: Seasonal Climate Outlooks using SCOPIC for
May to July 2015**

Predictors and Period used: Nino 3.4 values from January to March 2015

Station	Below Median (prob)	Median Rainfall (mm)	Above Median (prob)		LEPS	Hit-rate
Afiamalu	59	738.3	41		4.9%	57.6%
Nafanua	54	523.6	46		-2.5%	44.4%
Apia	57	410.2	43		3.6%	54.55
Faleolo	66	330.4	34		13.4%	67.9%

Station	Below Normal (prob)	33%ile rainfall (mm)	Normal (prob)	66%ile rainfall (mm)	Above Normal (prob)	LEPS	Hit-rate
Afiamalu	46	621.0	30	839.0	24	13.1%	39.4%
Nafanua	39	452.1	30	586.6	31	-1.6%	14.8%
Apia	37	331.7	33	480.8	30	-2.4%	30.3%
Faleolo	44	274.6	35	390.9	21	8.9%	50.0%

**TABLE 4: Seasonal Climate Outlooks using POAMA2 for
May to July 2015**

Station	Lower Tercile (prob)	33%ile rainfall (mm)	Middle Tercile (prob)	66%ile rainfall (mm)	Upper Tercile (prob)		
Apia	27	388	55	530	18		

Summary Statements

Rainfall for March 2015:

All stations recorded '**normal**' rainfall, except Faleolo which received 'above normal' rainfall.

Accumulated rainfall for January to March 2015, including outlook verification:

All stations received '**normal**' accumulated rainfall except Faleolo station which received 'below normal' rainfall.

Faleolo station was '**consistent**' with the outlook whereas Nafanua, Afiamalu and Apia were '**near consistent**'.

Outlooks for May to July 2015:

1. SCOPIC:

- '**Below normal**' rainfall is the most likely outcome for Afiamalu and Faleolo stations, with '**normal**' the next most likely.
- The outlook for Nafanua and Apia station give little guidance for the coming season as the chances of '**above normal**', '**normal**' and '**below normal**' rainfall are similar.
- The confidence of the model ranges from '**good**' to '**very low**'.

2. POAMA:

- '**Normal**' rainfall is favoured for Apia in the coming season.

NB: The X LEPS % score has been categorised as follows:

Very Low: $X < 0.0$

Low: $0 \leq X < 5$

Moderate $5 \leq X < 10$

Good: $10 \leq X < 15$

High: $15 \leq X < 25$

Very High: $25 \leq X < 35$

Exceptional: $X \geq 35$